

# Origin of the Neoproterozoic Baijianshan banded iron formation at the southeastern margin of the Tarim Block in NW China: implication for an extremely reducing ocean

Xiao-Shu Hao Chuan-Lin Zhang\* Hui-Chao Zhang Teng Ding Xian-Tao Ye

*College of Oceanography, Hohai University, Nanjing 210098, P.R. China*

**Abstract:** The Neoproterozoic Banded iron formations (BIFs) were closely associated with the “Snowball Earth” during the breakup of the Rodinia, thus they played an important role in our understanding of the atmospheric and oceanic oxygen levels during this period. In this contribution, the Neoproterozoic (ca. 737 Ma) Baijianshan BIF at Southeast Tarim, northwestern China was identified. Magnetite is the dominated iron-species, which occurs as the lamina interbedded with chert. The BIF contains low concentrations of trace elements, and is depleted in light rare earth elements (LREEs) based on comparison with the Post Archean Australian Shale (PAAS). In addition, the BIF exhibits slightly positive La-Eu anomalies, negligible Ce anomalies, insignificant Y anomalies, chondritic Y/Ho ratios (23-32), and slightly chondritic initial  $\epsilon_{\text{Nd}}$  ( $t = 737$  Ma) values (-0.45 to 1.46, averaging 0.37). All these features indicate that the precipitation of Baijianshan BIF was closely related to the submarine low-T hydrothermal fluids with little detrital contribution. Moreover, the Baijianshan BIF is characterized by the significant enrichment of heavy Fe isotopes, with  $\delta^{57}\text{Fe}_{\text{IRMM-014}}$  values ranging from 1.78 to 3.05‰, revealing the partial oxidation of  $\text{Fe}^{2+}$  into  $\text{Fe}^{3+}$  during the precipitation of this BIF. Our data suggest that the formation of Baijianshan BIF was closely associated with a significantly reducing ocean, which most likely was isolated from the oxidized atmosphere by a local ice sheet. This Neoproterozoic Baijianshan ocean has the initial oxygen levels as low as, or even lower than that of Archean and Paleoproterozoic oceans.

**Key words:** Neoproterozoic, banded iron formations (BIFs), Sr-Nd isotope, Fe isotope, Tarim block

\*Corresponding Author: E-Mail: zhangchuanlin@hhu.edu.cn

## 1 Introduction

Banded iron formations (BIFs), one of the most important iron sources in the world, are typical marine sedimentary rocks mainly precipitated during the Precambrian. BIF was firstly defined as a type of sedimentary rock containing more than 15% iron (James, 1954, 1966), which was later revised as thinly layered or laminated rock in which chert (or its metamorphic equivalent) alternating with layers that are composed mainly of iron minerals with the iron content ranging from 20% to 35% (James, 1983). The oldest BIFs precipitated at 3.8 Ga (Isua, Greenland), and then a large number of BIFs formed during the Archean-Paleoproterozoic (3.5-1.8 Ga), with a peak at ca. 2.5 Ga (Klein, 2005 and the references therein). They disappeared at ca. 1.8 Ga, and reoccurred in the Neoproterozoic (0.8-0.6 Ga) after a billion-year hiatus (Klein, 2005). It is commonly accepted that the precipitations of BIFs were closely related to fundamental changes in atmospheric and/or oceanic oxygen levels (i.e., Great Oxidation Event, GOE), although some key issues, including the source of the hydrothermal fluids, oxygen level and tectonic settings, are still under debate (Holland, 2002). In line with tectonic backgrounds, the BIFs can be subdivided into Algoma- and Superior-type (Gross, 1980), the Algoma-type generally precipitated nearby the volcano craters, in the sequences mainly composed of volcanic rocks, greywackes and turbidites (Klein and Beukes, 1992; Li et al., 2014; Sial et al., 2015), while the Superior-type were usually formed with sedimentary rock in shallow marine environment, such as passive continental margin (Gross and Mcleod, 1980).

Although the resources of Neoproterozoic BIFs are even much less than those of Archean and Paleoproterozoic, they are widely distributed worldwide, such as Rapitan BIF in Canada (Halverson et al., 2011), Wadi Karim and Um Anab BIF in Egypt (Basta et al., 2011), Sawawin BIF in Saudi Arabia (Stern et al., 2013), Jacadigo in Brazil (Freitas et al., 2011), Holowilena and Oraparinna in South Australia (Cox et al., 2016), the Kingston Peak Formation in California (Lechte et al., 2018) and Wadi Hamama in the Arabian-Nubian Shield (El-Rahman et al., 2020). They were also discovered in the Precambrian massifs in Phanerozoic orogenic belts of South China, such as Sanjiang in Guangxi Province (Yan et al., 2010), Shilu in Hainan Province (Xu et al., 2014), Xinyu in Jiangxi Province (Li et al., 2014; Shen et al., 2018; Zhu et al., 2019), the Shalong in the Central Tianshan (Lei et al., 2016, 2018) and Dahongliutan BIF in the Western Kunlun orogenic belt (Hu et al., 2017, 2020). Several

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/1755-6724.14801](https://doi.org/10.1111/1755-6724.14801).

models have been proposed for interpreting the formation of the Neoproterozoic BIFs: (1) During the break-up of Rodinia supercontinent, the increased magmatic activity and the formation of “sedimentary exhalation rift” induced the addition of Fe flux from submarine exhalative (Basta et al., 2011; Cox et al., 2013); (2) The thick ice sheet formed by “Snowball Earth” event isolated the hydrosphere and oxidized atmosphere, which led to the reduction of ocean and the dissolution of a large amount of Fe element. After the ice cover melted, the Fe was oxidized and precipitated into BIF as a result of the contact between ocean and atmosphere (Hoffman et al., 1998; Hoffman and Schrag, 2002; Klein and Ladeira, 2004; Halverson et al., 2011) and (3) during the late Neoproterozoic, there was a reduced S-poor and Fe-rich environment, which was favorable for Fe element existing as divalent ions in seawater, and then to be oxidized and precipitated (Johnston et al., 2010).

In this study, we report a newly identified Neoproterozoic Baijianshan BIF at the southeastern margin of the Tarim Block in NW China. The precipitation mechanism and the oceanic redox condition of the Baijianshan BIF was investigated based on detailed field observations coupled with systematic elemental and isotopic compositions analysis. This study sheds a new light on our understanding of the possible coupling of the Cryogenian snowball Earth event and the formation of Neoproterozoic BIFs.

## 2 Regional Geology

The Tarim Craton in NW China, covering an area of more than 600,000 km<sup>2</sup>, is one of the main three Precambrian nuclei in China (i.e., North China, South China and Tarim). It is surrounded by the Phanerozoic orogenic belts of Tianshan Mountains to the north, the western Kunlun Mountains to the south, and the Central-Southern Altyn Tagh Mountains to the southeast. Despite most of the Tarim Craton being covered by aeolian sands and difficulties in accessibility, previous work reveals that the craton is characterized by a typical double-layered structure consisting of a Pre-Cryogenian basement and Cryogenian-Cambrian cover sequences (Xinjiang BGMR, 1993; Zhang et al., 2013). Particularly, recent studies demonstrate that the craton docked at the northern fringe of the Rodinia slightly earlier than 760 Ma and then the Cryogenian-Cambrian sedimentary sequences deposited during the breakup of the Rodinia (Zhang et al., 2010, 2016; Xu et al., 2013).

Neoproterozoic sequences in Tarim mainly outcrop at its marginal areas such as the Aksu, Quruqtagh, Tiekelike and Altyn (Fig. 1a). Drilling boreholes interior of Tarim also revealed their equivalent sequences in the Central Tarim (Xu et al., 2013). Among them, tillite sequences were observed in Aksu, Quruqtagh and Tiekelike areas. Geochronological data demonstrated three phases of glaciation, i.e., ca. 740 Ma, ca. 635 Ma and ca. 580 Ma, were developed in the most well-preserved Cryogenian system in the Quruqtagh area (Xu et al., 2009). The two phases of tillites in Aksu were probably equivalent with two later phases of the tillite in Quruqtagh (Xu et al., 2013). As for the tillite in the Tiekelike area, despite that the detrital zircon U-Pb ages constrained its lower deposition age at ca. 740 Ma, its precise age is still unknown (Zhang et al., 2016).

The Neoproterozoic sequence in the Altyn is distributed in W-E direction along the southern margin of the early Precambrian Aktas terrane. It contacts with the Ordovician volcanic-sedimentary sequence (locally known as the Lapeiquan Formation) and the early Precambrian basement by faults (Zhang et al., 2014). In Chinese literatures it was suggested to be Mesoproterozoic and termed as Zhuoabulake Formation (Xinjiang BGMR, 1993) (Fig. 1b). Field observations reveal that the Zhuoabulake Formation can be subdivided into three members from the bottom to top (Fig. 2). The lower member is mainly composed of interlayered low-matured sandstone (greywacke) with minor slate, silicalite and volcanic debris with a five-meter-thick layer of conglomerate at its bottom. The middle member was composed of phyllite, slate and silicalite. Four layers of BIF, which is the focal point of the contribution, mainly comprised by silicalite and silicalite-magnetite, occur in this member (Fig. 3a-b). Additionally, three thin tuff layers (Fig. 3a) with the thickness of less than 30 centimeter, and one-meter-thick basalt layer, of one meter in thickness, occur at the lower part of the BIF layers. The upper member was mainly composed of black shale with minor slate. No tillite was observed in the Formation.

## 3 Petrographic Features

The iron ores are composed of interbedded magnetite-rich and quartz-rich lamina. The detailed petrographic features were examined by microscopes and scanning electron microscopy (SEM) (Fig. 3c-h).

Four layers of Fe orebodies exhibit visibly laminated and local bending features (Fig. 3b). The hand specimens show obvious stratabound features (Fig. 3c). Under the observation of SEM, most of the samples are well-banded with clear contacts between quartz-rich bands and magnetite-rich bands (Fig. 3d). Some “self- fragmentary structures” occur locally with the blurred contacts, which may be attributed to co-precipitation disturbance.

Eighteen BIF samples were collected from bottom to top of the four layers of Fe-ore bodies (Fig. 2).

These samples are dominated by magnetite and quartz with banded structures and mainly show mesoband (mm- to cm-scale) and microband (sub-mm- to mm-scale), which are composed of magnetite, quartz, rare hematite, siderite, albite and biotite (Fig. 3e-f). Quartz and other gangue minerals, such as chlorite (Fig. 3g) and sericite (Fig. 3h), occur occasionally as veinlets cutting across the bedding.

The shallow gray tuffs under the four layers of Fe ore bodies have been altered into clays. Nevertheless, the sub-euhedral or euhedral plagioclase and quartz crystal fragments, mostly less than 2 mm, can be seen under magnifier. The crystal fragments account for about 15-20% in the tuffs. One sample was collected from the tuff to constrain the deposition age of the Baijianshan BIF (N39°08'59.54", E91°50'38.90": 2071).

The blackish green basalt layer occurs at the bottom of the BIF layers, with the thickness of 1 meter. The minerals in the basalt were intensively altered and needle-like albite, chlorite as well as Ti-Fe oxide can be observed in thin sections, three samples were collected from the basalt layer for geochemical analysis (2702H1, 270H2 and 2702H3).

## 4 Analytical Methods

### 4.1 In situ zircon U-Pb isotope analysis

Conventional magnetic and density techniques were used to concentrate non-magnetic, heavy fractions and then zircon grains were hand-picked under a binocular microscope. After being mounted in epoxy mount, zircon grains were then polished to section the crystals in half for analysis. All zircons were documented with transmitted and reflected light micrographs as well as cathodoluminescence (CL) images to reveal their internal structures. Zircon U-Pb ages were analyzed using the LA-ICP-MS at the Tianjin Institute of Geology and Mineral Resources (China Geological Survey). Detailed analytical procedures can be found in Li et al. (2009). Zircon standards 91500 and GJ-1 (Jackson et al., 2004) were used to monitor the analysis. The U-Pb concordia plots were processed by ISOPLOT 3.0 and data were presented with  $1\sigma$  errors and 95% confidence limits (Ludwig, 2003). The zircon U-Pb age data are listed in Supplementary Table 1.

### 4.2 Whole rock geochemistry

Seventeen BIF samples and three basalt samples were carefully selected along the geological location. Whole-rock major compositions were analyzed using standard X-ray fluorescence (XRF) on fused glass beads at the Nanjing Institute of Geology and Mineral Resources (China Geological Survey), following the procedures similar to those described by Li et al., (2006a, b). Analytical uncertainties are between 1 and 5%. In addition, we used wet chemistry technique to measure FeO and Fe<sub>2</sub>O<sub>3</sub>, with the procedures described by Andrade et al. (2002). The measured data are listed in supplementary Table 2.

Trace elements were determined using a Perkin-Elmer Sciex ELAN DRC-e ICP-MS at the State Key Laboratory of Ore Deposit Geochemistry, Institute of Geochemistry (Chinese Academy of Sciences), with the analytical accuracy better than 5%. Samples were digested with 1ml of HF and 0.5 ml of HNO<sub>3</sub> in screw top PTFE-lined stainless steel bombs at 190°C for 12h (Qi et al., 2000). The analytical precision for most elements was better than 1% with concentrations > 200 ppm, and 1-3% when less than 200 ppm. The analytical results are reported in supplementary Table 2.

Sr-Nd isotopes were measured using the Micromass Isoprobe Multi-collector MC-ICP-MS at Tianjin Institute of Geology and Mineral Resources (China Geological Survey), with the analytical procedures similar to those reported by Li et al. (2004). Measured <sup>87</sup>Sr/<sup>86</sup>Sr and <sup>143</sup>Nd/<sup>144</sup>Nd ratios were corrected for mass-fractionation using <sup>86</sup>Sr/<sup>88</sup>Sr = 0.1194 and <sup>146</sup>Nd/<sup>144</sup>Nd = 0.7219, respectively. The measured values for the NBS SRM 987 standard <sup>87</sup>Sr/<sup>86</sup>Sr = 0.71025 and the Shin Etsu JNdi-1 standard <sup>143</sup>Nd/<sup>144</sup>Nd = 0.512115. All the initial <sup>143</sup>Nd/<sup>144</sup>Nd values of the BIF are calculated with a probable depositional age (737 Ma) for the Baijianshan BIF.

Iron isotope ratios were measured using MC-ICP-MS at Beijing Createch Testing Technology Co., Ltd. The results of Fe isotope ratio determination as the ten thousandth deviation of the sample from the standard sample is as follows:

$$\delta^x Fe/^{54}Fe(\text{‰}) = \left[ \frac{({}^x Fe/^{54} Fe)_{\text{sample}}}{({}^x Fe/^{54} Fe)_{\text{standard}}} - 1 \right] \times 1000$$

(x = 56, 57)

The performance of the instrument was assessed by repeated measurements of an in-house standard (CAGS Fe) that yielded deviations relative to the IRMM-014 Fe isotope reference material. The average Fe isotope values for CAGS Fe are  $\delta^{56}Fe_{\text{IRMM014}}\text{‰} = \delta^{56}Fe_{\text{CAGS}}\text{‰} + 0.80$  and  $\delta^{57}Fe_{\text{IRMM014}}\text{‰} = \delta^{57}Fe_{\text{CAGS}}\text{‰} + 1.20$ , using the procedures described by Tang et al. (2016).

## 5 Analytical Results

### 5.1 Zircon U-Pb age

Zircons from the tuff sample 2071 are variable in size with the length varying from 60  $\mu\text{m}$  to 150  $\mu\text{m}$  and the aspect ratios 1-2. In CL images most zircon exhibit oscillatory zoning, sharing the features of the zircons crystallized from silicic magma (Wu and Zheng, 2004). Thirty-two analyses were conducted on 32 zircon grains and the results are presented in supplementary Table 1 and illustrated in Fig. 4. Obviously, some analyses show variable radiogenetic lead lost. However, the results can be broadly divided into two sub-groups. Group 1, which is likely to be xenocrystal and generally have larger size, constructs a good Discordia with upper intercepted age of  $1378\pm 44$  Ma (MSWD=5.8). Group 2 mostly shows euhedral in form and smaller in size. Twenty-one analyses construct a good Discordia with an intercepted age of  $736.2\pm 3.8$  Ma (N=21, MSWD=1.08) (insert of Fig. 4). Excluded the significant discordant analyses, the left fifteen analyses of well concordance between  $^{206}\text{Pb}/^{238}\text{U}$  age and  $^{207}\text{Pb}/^{235}\text{U}$  age, yield a mean  $^{206}\text{Pb}/^{238}\text{U}$  age of  $737.0\pm 4.0$  Ma (N=15, MSWD=0.20). This age is interpreted as the deposition time of the Baijianshan BIF.

### 5.2 Whole rock Geochemistry

#### 5.2.1 Geochemistry of the basalts

Three basalt samples (2702H1, 270H2 and 2702H3) have low  $\text{SiO}_2$  contents (39.93% to 44.55%). The incompatible elements for the rock type classification are used due to the high LOI. Their high Nb/Y ratios (1.05-1.09) define their alkaline signature and in the Nb/Y vs. Zr/TiO<sub>2</sub> diagram, they plot into the alkaline basalt field (Figure not shown). With respect to the trace elements, they have  $\Sigma\text{REE}$  ranging from 186 ppm to 195 ppm and show variable LREE enriched Chondrite-normalized pattern (Fig. 5a) ( $\text{La}_\text{N}/\text{Yb}_\text{N} = 9.8\text{-}11.0$ ). In line with their low Cr (214-306 ppm) and Ni contents (162-216.9 ppm), the basalts are evolved magma. However, the primitive mantle-normalized diagram shares most features of the OIB-like basalts with insignificant Nb-Ta trough (Nb/La = 1.0-1.1) (Figure not shown).

#### 5.2.2 Geochemistry of the BIF

As shown in supplementary Table 2, the samples from Baijianshan BIF are rich in  $\text{Fe}_2\text{O}_3^\text{T}$  and  $\text{SiO}_2$ .  $\text{Fe}_2\text{O}_3^\text{T}$  concentrations vary between 13.11% and 56.65% (38.44% on average), whereas  $\text{SiO}_2$  varies between 26.81% and 55.06% (42.02% on average). They have low contents of  $\text{Al}_2\text{O}_3$  (1.56%-7.58%, 3.66% on average),  $\text{TiO}_2$ , MnO,  $\text{P}_2\text{O}_5$ , CaO,  $\text{Na}_2\text{O}$ ,  $\text{K}_2\text{O}$  and variable total rare earth elements and other incompatible elements (supplementary Table 2).

Their REE abundances vary from 38.1 ppm to 150.8 ppm. Normalized against Post Archean Average Shale (PAAS; McLennan, 1989), they are characterized by depletion of LREE relative to HREE with the  $\text{La}_\text{N}/\text{Yb}_\text{N}$  of 0.23-0.69 (Fig. 5b). The REY patterns exhibit no Ce and Pr anomalies (the combination of  $\text{Ce}/\text{Ce}^* \sim 1$  and  $\text{Pr}/\text{Pr}^* \sim 1$ ; Bau and Dulski, 1996) and insignificant Eu and Y anomaly ( $\text{Eu}/\text{Eu}^* = 0.96\text{-}1.14$ ; Y/Ho ratios ranging from 23.3 to 31.8 with an averaged value of 26.39, Table 2).

#### 5.2.3 Sr-Nd isotope compositions

Sr-Nd concentrations and isotopic ratios of the Baijianshan BIF and basalts are presented in supplementary Table 3. As for the basalts, the basalts exhibit high initial Sr isotope compositions with ( $^{87}\text{Sr}/^{86}\text{Sr}$ )<sub>i</sub> ranging from 0.7061 to 0.7084 due to post-eruption hydrothermal alteration. In addition, they have pronounced depleted Nd isotope compositions with  $\epsilon\text{Nd}$  (t = 737 Ma) ranging from 5.7 to 6.9.

Strontium and neodymium concentrations of Baijianshan BIF range from 20.3 ppm to 256 ppm and from 7.06 ppm to 59.8 ppm, respectively. The nine samples have a large range of measured  $^{143}\text{Nd}/^{144}\text{Nd}$  (0.51231-0.51243) and  $^{87}\text{Sr}/^{86}\text{Sr}$  (0.7179-0.7463) ratios. The  $\epsilon\text{Nd}$  (t = 737Ma) values range from -0.45 to 1.46, with an average of 0.37. However, the BIF samples exhibit a large range of the ( $^{87}\text{Sr}/^{86}\text{Sr}$ )<sub>i</sub> values between 0.6412 and 0.7175, possibly due to the post-deposition hydrothermal alteration and the large range of Rb/Sr ratios varying from 0.02 to 3.47.

#### 5.2.4 Fe isotope compositions

The Fe isotope compositions of the Baijianshan BIF samples are presented in supplementary Table 4. The six samples yielded  $\delta^{56}\text{Fe}_{\text{IRMM-014}}$  values of 1.24-2.17‰ and  $\delta^{57}\text{Fe}_{\text{IRMM-014}}$  values of 1.78-3.05‰. Their enrichment in heavy Fe isotopes shares the signatures of the Neoproterozoic BIFs (Halverson et al., 2011; Yan et al., 2011; Shen et al., 2018).

## 6 Discussions

### 6.1 Iron sources for the Baijianshan BIFs

Field and microscopic observations revealed that the Neoproterozoic Baijianshan BIF has not

experienced significant metamorphism (Fig. 3b, c and d), indicating the original geochemical characteristics were well retained and can be used to decipher the iron sources and the oceanic environments when it precipitated.

The Baijianshan BIF shows a modern seawater REE signature with significant LREE depletion relative to the HREE ( $La_N/Yb_N = 0.41$ ) (Fig. 5b, Zhang and Nozaki, 1996; Alibo and Nozaki, 1999). A range of factors can affect the primary REY composition in terms of BIF depositional processes, including post-depositional metamorphism and/or syn-deposition clastic contamination (Alexander et al., 2008; Hu et al., 2020). Previous studies demonstrated that diagenetic and metamorphic events (such as weathering and/or fluid-rock interactions) have negligible effects on the REY of the chemical sediments (Bolhar et al., 2004). However, Rb is a relatively mobile element compared to highly immobile elements such as Th. In Fig. 6a, the well positive correlation between Rb and Th demonstrates insignificant mobility of the LILEs (Large Ion Lithophile Elements) and the absence of significant diagenetic or metamorphic alteration effects, this conclusion is also consistent with thin section observations (Fig. 3c, d, g, h). With respect to the syn-depositional processes, the contents of  $Al_2O_3$  and  $TiO_2$  are consistently low with averages of 3.66% and 0.41%. In addition, despite the positive correlations between  $Al_2O_3$  and  $TiO_2$  ( $r = 0.93$ ),  $K_2O$  ( $r = 0.79$ ), Rb ( $r = 0.78$ ), V ( $r = 0.45$ ), Sc ( $r = 0.77$ ), Cs ( $r = 0.66$ ), Cr ( $r = 0.72$ ), and  $\sum REE$  ( $r = 0.62$ ) (Fig. 6b-i) signify the incorporation of terrigenous contribution in the deposition processes, most correlation coefficients are much lower than those BIFs sourced from recycled crust (Basta et al., 2011; Cox et al., 2013; Hu et al., 2017, 2020). Besides, the Baijianshan BIF shares similar REE patterns with those BIFs from North China Craton but contains slightly higher contents of  $Al_2O_3$  and  $TiO_2$  (Li et al., 2014). Taken together, we suggest a little incorporation of terrigenous component in the Baijianshan BIF.

Positive Eu anomalies are generally interpreted as the precipitation of high-temperature hydrothermal fluids (Danielson et al., 1992; Bau and Dulski, 1999) while low-temperature hydrothermal fluids usually display weak or no Eu anomalies (Michard et al., 1993; Li et al., 2014). Most samples from the Baijianshan BIF show negligible Eu anomalies ( $Eu/Eu^* = 0.96\sim 1.14$ , averaging 1.07), which are indicative of low-temperature hydrothermal solutions (Danielson et al., 1992). The  $\epsilon Nd(t)$  value can be used as a tracer to distinguish different sources and crustal contamination. As shown in Fig. 7, the Baijianshan BIF has slightly average positive  $\epsilon Nd(t)$  value (average of  $\epsilon Nd(t) = 0.37$ ), much lower than those of Baijianshan basalt, indicating that the deposition of the Baijianshan BIF was controlled by a hydrothermal flux with mantle-like positive  $\epsilon Nd(t)$  values (Jacobsen and Pimentel-Klose, 1988). However, the slightly negative correlation between  $\epsilon Nd$  (737 Ma) and  $Al_2O_3$  contents (Fig. 8a) argues for the low proportion of detrital components inputting.

The Y/Ho ratios of BIFs can provide constraints of the material source and precipitation environment of BIFs (Hu et al., 2017). Modern seawaters have a Y/Ho ratio of 44-74 (Bau and Dulski, 1996). The terrestrial material has a Y/Ho ratio of  $\sim 26$ , any little terrestrial contamination could quickly descend seawater-like superchondritic Y/Ho ratios ( $>44$ ) (Bolhar et al., 2004). The hydrothermal fluids have almost chondritic Y/Ho ratios (26-28, Douville et al., 1999; Bau and Dulski, 1999). The Baijianshan BIF has a range of Y/Ho ratios between 23 and 32, similar to the chondritic values (26-28, Bau and Dulski, 1999), possibly due to the mixture of high and/or low-T hydrothermal fluids (Hu et al., 2020). Y/Ho ratios of the Baijianshan BIF samples might inherit from the low-T hydrothermal fluids, but the influences of terrestrial materials can not be completely excluded. Compared with the BIFs in North China Craton, the Baijianshan BIF displays relatively low positive Eu anomalies, low Y/Ho ratios as well as a slightly depleted  $\epsilon Nd(t)$  values (Li et al., 2014; Wang et al., 2016), illustrating the input of low-temperature hydrothermal fluids during the deposition of the Baijianshan BIF.

To further ascertain the sources of the Baijianshan BIF, two-components mixing model invoked by Alexander et al. (2008) is used in the study. We consider that a mixture of hydrothermal fluid and seawater is sufficient for the production of observed Eu/Sm and Sm/Yb values of the Baijianshan BIF. As shown in Fig. 9, the Baijianshan BIF samples straddle between low-temperature hydrothermal fluid and seawater but far away from the high-temperature hydrothermal fluid, similar to the scenario of the Xinyu BIF. Therefore, we suggest that Baijianshan BIF is predominantly controlled by seawater and low-T hydrothermal solutions, accompanied with a minor amount of detritus input.

OIB-like geochemistry of the basalt layer (one meter in thickness) at the lower part of the BIF layers together with the sedimentary features of the BIF-bearing sequences, unambiguously demonstrated a rifting background in Tarim during the Cryogenian. At the same time, intensive magmatic activities could significantly enhance the submarine hydrothermal circulations, resulting in the increase of Fe fluxes into seawater. This conclusion is consistent with the multiple episodes of hydrothermal chimneys identified in nearby areas, such as South China and the Tarim Basin (Chen et al., 2009; Zhou et al., 2014). The dissolved Fe thus might be the main sources for the precipitation of the Baijianshan BIF.

## 6.2 Oceanic redox states during the precipitation of the Baijianshan BIF

Ce is sensitive to the redox environment of the seawater (Feng et al., 2016). In Fig. 8b, the consistent

Ce/Ce\*<sub>PAAAS</sub> values with variable Al<sub>2</sub>O<sub>3</sub> contents of the Baijianshan BIF indicate that terrigenous detrital materials played little role in the geochemical budget of Ce. In oxidized seawater, Ce(III) transforms into Ce(IV), and then Ce(IV) is likely to be hydrolyzed and precipitated with Fe-Mn oxyhydroxides, organics and clay which lead to a significant negative Ce anomaly of seawater (Byrne and Sholkovitz, 1996). As shown in the Ce/Ce\* vs. Pr/Pr\* diagram (Fig. 10), all the Baijianshan BIF samples display no Ce anomalies, similar with most Neoproterozoic BIFs, unambiguously arguing for anoxic environment in the ancient ocean. Although the atmospheric and oceanic oxygen levels during the Neoproterozoic are still under debate, numerous studies revealed that Neoproterozoic oceans underwent a stepwise and protracted oxidation. Anoxic ferruginous deep seawater was a typical feature of the late Neoproterozoic, as inferred from geochemical proxies such as iron geochemistry (e.g., Canfield et al., 2008; Sperling et al., 2015), redox-sensitive elements (e.g., Schröder and Grotzinger, 2007; Rajabi et al., 2015), and framboidal pyrite (e.g., Rajabi et al., 2015). Stern et al. (2013) suggested that this scenario might be attributed to the seawater surfaces being covered by ice sheets, blocking the oxygen from atmosphere dissolving into the widespread anoxic ferruginous ocean. Despite no tillite was identified at the Baijianshan BIF, the coeval tillite sequences were documented in the Quruqtagh area (i.e., ca.740 Ma Beiyixi tillite) of NE Tarim (Xu et al., 2009), which argued for the possibility of ice sheets in the Cryogenian Baijianshan ocean.

In a Fe<sup>2+</sup>-rich marine settings, both abiotic and biotic action were demonstrated as oxidation pathways for the precipitation of iron oxyhydroxides (Fe(OH)<sub>3</sub>) from dissolved Fe<sup>2+</sup> (Konhauser et al., 2011). As an element of variable valency, the fractionation of Fe isotope is affected by the oxidation-reduction condition (Bullen et al., 2001; Zhu et al., 2002; Balci et al., 2006). Experimental studies demonstrated that oxidation of Fe<sup>2+</sup> to Fe<sup>3+</sup> in solution causes considerable Fe isotope fraction, generating an enrichment of heavy Fe isotopes in Fe<sup>3+</sup> (Bullen et al., 2001; Johnson et al., 2002; Balci et al., 2006; Hou et al., 2014). On the other hand, the fractionation caused by pyrite with light-Fe isotopes is considered to play a leading role in the enrichment of heavy Fe isotopes in the residual system relative to the original state (Rouxel et al., 2016), since the pyrite is easier to precipitation from plume. Nevertheless, there is no sulfide observed in the Baijianshan BIF, thus the potential influences of sulfides on the enrichment of heavy Fe isotopes can be excluded. Under oxidized environment, the enrichment of heavy Fe isotopes is shown in oxide or hydroxide of iron trivalent (Fe<sup>3+</sup><sub>ppt</sub>) and light Fe isotope is displayed in Ferrous solution (Fe<sup>2+</sup><sub>aq</sub>) (Johnson et al., 2002). The magnitude of Fe isotope fractionation is controlled by the degree of precipitation from Fe<sup>2+</sup> to Fe<sup>3+</sup>, which is related to the degree of ocean oxidation (Yan et al., 2010). The Fe in the seawater can be completely precipitated and no fractionation of the Fe isotope occurs when the seawater is completely oxidized, thus the Fe isotope values in iron oxide precipitates (Fe<sup>3+</sup>) can represent the Fe isotope information of seawater. However, when the seawater is partially oxidized, the Fe in the seawater is partially precipitated and the Fe isotope fractionation will occur in iron oxide precipitates (Fe<sup>3+</sup>). Therefore, the Fe isotope value can be used as a proxy to decipher the redox state of seawater (Yan et al., 2010). Compared with some BIFs deposited during the Archean-Paleoproterozoic (Dauphas et al., 2004; Rouxel et al., 2005; Li et al., 2012; Johnson et al., 2003; Hou et al., 2014; Li et al., 2014), the Neoproterozoic Baijianshan, Sanjiang (Yan et al., 2010) and Xinyu BIFs (Shen et al., 2008) have higher values in δ<sup>57</sup>Fe (Fig. 11). Several studies demonstrated that the low δ<sup>57</sup>Fe values of BIFs could be due to the contributions from a continental component and the continental Fe source is best explained by Fe mobilization on the continental margin by microbial dissimilatory iron reduction (DIR) (Johnson et al., 2008; Li et al., 2015). As for the Fe isotope values of Archean-Paleoproterozoic BIFs we cited, the Fe source of SW Greenland BIF, Anshan-Benxi BIFs, Gongchangling BIFs and Yuanjiacun BIFs are unambiguously demonstrated to be hydrothermal type similar to the scenario of the Baijianshan BIF (Dauphas et al., 2004; Li et al., 2012; Li et al., 2014; Hou et al., 2014), while the Fe of Transvaal, Manjeri, Belingwe Belt and Zimbabwe BIFs are probably from multiple sources (abiologic and biologic processes) (Johnson et al., 2008; Grassineau et al., 2011). The comparison reveals that the environment in which they precipitated was more reduced. Both the iron isotope and elemental geochemistry thus demonstrate that the Neoproterozoic Baijianshan ocean might have the same reduced environment as Archean or Paleoproterozoic, or even more reduced.

### 6.3 A model of the Baijianshan BIF

During the late Neoproterozoic period (i.e., 750-550 Ma), the Earth experienced long-lived global glaciations, known as the Snowball Earth events (Hoffman et al., 1998). In view of the Neoproterozoic oxygenation event (NOE) at that time, the content of oxygen in atmosphere was much higher than before (Och et al., 2012). Recently, EI-Rahman et al. (2020) argued that some Neoproterozoic BIFs are not glaciogenic due to their significant depleted Nd isotope compositions. Under this scenario, the iron was most likely deriving from hydrothermal alteration of juvenile oceanic crust. The Chondritic-like Nd isotope compositions of the Baijianshan BIF (average of εNd(t) = 0.37) do not favor this model. Furthermore, the close temporally-spatially related tillite sequences in the Quruqtagh of NE Tarim (Xu et al., 2009), 765 Ma-735 Ma tillites in Namibia, suggested a local glaciation termed as Kaigas

glaciation (Kay et al., 2001; Cailteux et al., 2005; Hoffmann et al., 2006). Considering the high latitude position of the Tarim in the Rodinia configuration (Li et al., 2008), the Kaigas glaciation (corresponding to the Beiyixi tillite in Tarim) most possibly occurred in Tarim. In combination with previous studies on other BIFs with the data presented in this study, we construct a cartoon model showing the formation mechanism of the Baijianshan BIF (Fig. 12).

Stage 1, the seawater was covered by ice sheet which led to the insulation between atmosphere and oceans with the influence of the “Snowball Earth”. And then it induced the emergence of a reductive ocean. The iron supplied by low-temperature hydrothermal fluids could exist as the ferrous form, forming a relatively stable anoxic ferruginous reservoir in the ocean.

Stage 2, at interglacial period, the partial melting of the ice sheet caused some oxygen from the atmosphere into the ocean, which emerged a stratified seawater. There were mainly anoxic and Fe<sup>2+</sup>-rich deeper seawater, whereas the Fe<sup>2+</sup> was gradually oxidized into Fe<sup>3+</sup> and precipitated into iron oxide in near-shore oxic shallow seawaters. The low-temperature hydrothermal was the dominant source during the precipitation process to interpret the slight positive Eu anomalies and positive values of εNd(t). Under the effect of the melting ice-sheet, the increasing weathering then contributed to a small amount of terrestrial materials into the shallow seawater, resulting the precipitation of the Baijianshan BIF.

## 7 Conclusions

The Baijianshan BIF precipitated in glacially-influenced settings during the Neoproterozoic (737 Ma). Low-T hydrothermal fluids, accompanied with limited detritus input, were the dominated iron sources for the formation of Baijianshan BIF.

The Baijianshan ocean was initially covered by ice sheets, giving rise to the extremely reduced environment favored by Fe<sup>2+</sup>. Then it was oxidized into Fe<sup>3+</sup> and precipitated as magnetite due to the thawing of ice sheets. At the early stage, local reduced ocean enhanced Fe fluxes genetically related to the intensive marine magmatism and hydrothermal activities due to the existence of ice sheets. Thawing of ice sheets induced oxidized oceanic layer, favoring the formation of the Neoproterozoic BIF at the later stage.

## Acknowledgements

We sincerely thank the anonymous reviewers for their careful and constructive comments on our manuscript, which significantly elevate the quality of this study. We are grateful to Prof. Hai-Bo Zou, for his careful polishing on the English writing. This project is funded by the Fundamental Research Fund for Central Universities (B16020127).

## References

- Alexander, B.W., Bau, M., Andersson, P., Dulski, P., 2008. Continentally-derived solutes in shallow Archean seawater: rare earth element and Nd isotope evidence in iron formation from the 2.9 Ga Pongola Supergroup, South Africa. *Geochim. Cosmochim. Acta* 72, 378-394.
- Alibo, D.S., Nozaki, Y., 1999. Rare earth elements in seawater: particle association, shale-normalization and Ce oxidation. *Geochim. Cosmochim. Acta* 63(3-4), 363-372.
- Andrade, S., Hypolito, R., Ulbrich, H.H.G.J., Silva, M.L., 2002. Iron(II) oxide determination in rocks and minerals. *Chem. Geol.* 182, 85-89.
- Balci, N., Bullen, T.D., Witte-Lien, K., Shanks, W.C., Motelica, M., Mandernack, K.W., 2006. Iron isotope fractionation during microbially stimulated Fe(II) oxidation and Fe(III) precipitation. *Geochim Cosmochim Acta* 70, 622-639.
- Basta, F.F., Maurice, A.E., Fontboté, L., Favarger, P.Y., 2011. Petrology and geochemistry of the banded iron formation (BIF) of Wadi Karim and UmAnab, Eastern Desert, Egypt: implications for the origin of Neoproterozoic BIF. *Precambrian Res.* 187, 277-292.
- Bau, M., Dulski, P., 1996. Distribution of yttrium and rare-earth elements in the Penge and Kuruman iron-formations, Transvaal Super group, South Africa. *Precambrian Res.* 79, 37-55.
- Bau, M., Dulski, P., 1999. Comparing yttrium and rare earths in hydrothermal fluids from the Mid-Atlantic Ridge: implications for Y and REE behaviour during nearvent mixing and for the Y/Ho ratio of Proterozoic seawater. *Chem. Geol.* 155, 77-90.
- Bolhar, R., Kamber, B.S., Moorbath, S., Fedo, C.M., Whitehouse, M.J., 2004. Characterisation of early Archaean chemical sediments by trace element signatures. *Earth Planet. Sci. Lett.* 222, 43-60.
- Bullen, T.D., White, A.F., Childs, C.W., Vivit, D.V., Schulz, M.S., 2001. Demonstration of significant abiotic iron isotope fractionation. *Geology* 29, 699-702.
- Byrne, R., Sholkovitz, 1996. Marine chemistry and geochemistry of the lanthanides. In: Gschneider Jr.K.A., Eyring, L. (Eds.), *Handbook on the Physics and Chemistry of the Rare Earths*. Elsevier, Amsterdam.
- Cailteux, J.L.H., Kampunzu, A.B.H., Batumike, M.J., 2005. Lithostratigraphic position and petrographic characteristics of R. A. T. (“Roches Argilo-Talqueuses”) Subgroup, Neoproterozoic Katangan Belt (Congo). *Journal of African Earth Sciences* 42, 82-94.
- Canfield, D.E., Poulton, S.W., Knoll, A.H., Narbonne, G.M., Ross, G., Goldberg, T., Strauss, H., 2008. Ferruginous Conditions Dominated Later Neoproterozoic Deep-Water Chemistry. *Science* 321(5891), 949-952.
- Chen, D.Z., Wang, J.G., Qing, H.R., Yan, D.T., Li, R.W., 2009. Hydrothermal venting activities in the Early

- Cambrian, South China: Petrological, geochronological and stable isotopic constraints. *Chemical Geology* 258, 168-181.
- Cox, G.M., Halverson, G.P., Minarik, W.G., Heron, D.P.L., Macdonald, F.A., Bellefroid, E.J., Strauss, J.V., 2013. Neoproterozoic iron formation: an evaluation of its temporal, environmental and tectonic significance. *Chem. Geol.* 362, 232-249.
- Cox, G.M., Halverson, G.P., Poirier, A., Le Heron, D., Strauss, J.V., Stevenson, R., 2016. A model for Cryogenian iron formation. *Earth Planet. Sci. Lett.* 433, 280-292.
- Danielson, A., Moller, P., Dulski, P., 1992. The europium anomalies in banded iron formations and the thermal history of the oceanic crust. *Chem. Geol.* 97, 89-100.
- Dauphas, N., van Zuilen, M., Wadhwa, M., Davis, A.M., Marty, B., Janney, P.E., 2004. Clues from Fe isotope variations on the origin of early Archean BIFs from Greenland. *Science* 306, 2077-2080.
- EI-Rahman, Y.A., Gutzmer, J., Li, X.H., Seifert, T., Li, C.F., Ling, X.X., Li, J., 2020. Not all Neoproterozoic iron formations are glaciogenic: Sturtian-aged non-Rapitan exhalative iron formations from the Arabian-Nubian Shield. *Mineralium Deposita* 55, 577-596.
- Feng, L.J., Huang, J., Lu, D.B., Zhang, Q.R., 2016. Major and trace element geochemistry of the Neoproterozoic syn-glacial Fulu iron formation, South China. *Geological Magazine* 154, 1-10.
- Freitas, B.T., Warren, L.V., Boggiani, P.C., Almeida, R.P.D., Piacentini, T., 2011. Tectono-sedimentary evolution of the Neoproterozoic BIF-bearing Jacadigo Group, SW-Brazil. *Sedimentary Geology* 238, 48-70.
- Grassineau, N.V., Nisbet, E.G., Bickle, M.J., Fowler, C.M.R., Lowry, D., Matthey, D.P., Abell, P., Martin, A., 2000. Antiquity of the biological sulphur cycle: evidence from sulphur and carbon isotopes in 2700 million-year-old rocks of the Belingwe Belt, Zimbabwe. *Proc. R. Soc. Lond. B* 268, 113-119.
- Gross, G.A., 1980. A classification of iron-formation based on depositional environments. *Canadian Mineralogist* 18, 215-222.
- Gross, G.A., Mcleod, C.R., 1980. A preliminary assessment of the chemical composition of iron formation in Canada. *Canadian Mineralogist* 18, 223-229.
- Halverson, G.P., Poitrasson, F., Hoffman, P.F., Nédélec, A., Montel, J.M., Kirby, J., 2011. Fe isotope and trace element geochemistry of the Neoproterozoic syn-glacial Rapitan iron formation. *Earth Planet. Sci. Lett.* 309(1), 100-112.
- Hoffman, P.F., Kaufman, A.J., Halverson, G.P., Schrag, D.P., 1998. A Neoproterozoic snowball Earth. *Science* 281, 1342-1346.
- Hoffman, P.F., Schrag, D.P., 2002. The snowball earth hypothesis: testing the limits of global change. *Terra Nova* 14(3), 129-155.
- Hoffmann, K.H., Condon, D.J., Bowring, S.A., Prave, A.R., Fallick, A., 2006. Lithostratigraphic, carbon ( $\delta^{13}\text{C}$ ) isotope and U-Pb zircon age constraints on early Neoproterozoic (ca. 755 Ma) glaciation in the Gariep Belt, southern Namibia. *Snowball Earth Conference*, July 16-21, 2006, Montle Namibia, Ticino, Switzerland, 51.
- Holland, H.D., 2002. Volcanic gases, black smokers, and the great oxidation event. *Geochimica et Cosmochimica Acta* 66(21), 3811-3826.
- Hou, K.J., Li, Y.H., Gao, J.F., Liu, F., Qin, Y., 2014. Geochemistry and Si-O-Fe isotope constraint on the origin of banded iron formations of the Yuanjiacun Formation, Lvliang Group, Shanxi, China. *Ore Geology Reviews* 57, 288-298.
- Hu, J., Wang, H., Wang, M., 2017. Geochemistry and origin of the Neoproterozoic Dahongliutan banded iron formation (BIF) in the Western Kunlun orogenic belt, Xinjiang (NW China). *Ore Geology Reviews* 89, 836-857.
- Hu, J., Wang, H., Zhang, L.G., 2020. A rare earth element and Nd isotopic investigation into the provenance and deposition of the Dahongliutan banded iron formation and associated carbonates, NW China: Implications on Neoproterozoic seawater compositions. *Precambrian Research* 342, 1-29.
- Jackson, S.E., Pearson, N.J., Griffin, W.L., Belousova, E.A., 2004. The application of laser ablation-inductively coupled plasma-mass spectrometry to in situ U-Pb zircon geochronology. *Chemical Geology* 211(1-2), 47-69.
- Jacobsen, S.B., Pimentel-Klose, M.R., 1988. Nd isotopic variations in Precambrian banded iron formations. *Geophys Res Lett* 15, 393-396.
- James, H.L., 1954. Sedimentary facies of iron-formation. *Economic Geology* 49(3), 235-293.
- James, H.L., 1966. Chemistry of the iron-rich sedimentary rocks. *U.S. Geol. Survey Prof.* 440-461.
- James, H.L., 1983. Distribution of banded iron-formation in space and time. *Developments in Precambrian Geology* 6, 471-490.
- Johnson, C.M., Beard, B.L., Beukes, N.J., Klein, C., O'Leary, J.M., 2003. Ancient geochemical cycling in the Earth as inferred from Fe isotope studies of banded iron formations from the Transvaal Craton. *Contributions to Mineralogy and Petrology* 144(5), 523-547.
- Johnson, C.M., Beard, B.L., Klein, C., Beukes, N.J., Roden, E.E., 2008. Iron isotopes constrain biologic and abiologic processes in banded iron formation genesis. *Geochimica Et Cosmochimica Acta* 72(1), 151-169.
- Johnson, C.M., Skulan, J.L., Beard, B.L., Sun, H., Neelson, K.H., Braterman, P.S., 2002. Isotopic fractionation between Fe(III) and Fe(II) in aqueous solution. *Earth Planet Sci Lett* 195, 141-153.
- Johnston, D.T., Poulton, S.W., Dehler, C., Porter, S., Husson, J., Canfield, D.E., Knoll, A.H., 2010. An emerging picture of Neoproterozoic ocean chemistry: insights from the Chuar Group, Grand Canyon, USA. *Earth Planet. Sci. Lett.* 290, 64-73.
- Kay, R.M., Liyungu, A.K., Njamu, F.M., Somwe, V., Banda, J., Mosley, P.N., Armstrong, R.A., 2001. The western arm of the Lufilian Arc in NW Zambia and its potential for copper mineralization. *Journal of African Earth Sciences* 33, 503-528.
- Klein, C., 2005. Some Precambrian banded iron-formations (BIFs) from around the world: Their age, geologic setting, mineralogy, metamorphism, geochemistry, and origins. *American Mineralogist* 90(10), 1473-1499.
- Klein, C., Beukes, N.J., 1992. Time distribution, stratigraphy, sedimentologic setting, and geochemistry of Precambrian iron-formations. In: Schopf, J.W., Klein, C. (Eds.), *The Proterozoic Biosphere*. Cambridge University Press, Cambridge, 139-146.
- Klein, C., Ladeira, E.A., 2004. Geochemistry and mineralogy of Neoproterozoic banded iron-formations and some selected, siliceous manganese formations from the Urucum district, Mato Grosso do Sul, Brazil. *Econ. Geol.* 99, 1233-1244.
- Konhauser, K.O., Kappler, A.K., Roden, E.E., 2011. The microbial role in iron redox and biomineralization reactions. *Elements* 7, 89-93.



- Lechte, M.A., Wallace, M.W., Hood, A. van S., Planavsky, N., 2018. Cryogenian iron formations in the glaciogenic Kingston Peak Formation, California. *Precamb. Res.* 310, 443-462.
- Lei, R.X., Wu, C.Z., Li, G.R., 2016. Neoproterozoic banded iron formation in the Central Tianshan, NW China: The Shalong example. *Acta Geologica Sinica (English Edition)* 90(5), 1543-1544.
- Lei, R.X., Wu, C.Z., Feng, Y.G., Xia, M.Z., Jiao, J.G., 2018. Formation age and geodynamic setting of the Neoproterozoic Shalong iron formation in the Central Tianshan, NW China: Constraints from zircon U-Pb dating, geochemistry, and Hf-Nd isotopes of the host rocks. *Geological Journal* 1-17.
- Li, H.M., Zhang, Z.J., Li, L.X., Zhang, Z.C., Chen, J., Yao, T., 2014. Types and general characteristics of the BIF-related iron deposits in China. *Ore Geology Reviews* 57, 264-287.
- Li, W.Q., Beard, B.L., Johnson, C.M., 2015. Biologically recycled continental iron is a major component in banded iron formations. *Proc. Natl. Acad. Sci.* 112, 8193-8198.
- Li, X.H., Liu, D.Y., Sun, M., Li, W.X., Liang, X.R., Liu, Y., 2004. Precise Sm-Nd and U-Pb isotopic dating of the super-giant Shizhuyuan polymetallic deposit and its host granite; Southeast China. *Geol. Mag.* 141, 225-231.
- Li, X.H., Li, Z.X., Wingate, M.T.D., Chung, S.L., Liu, Y., Lin, G.C., Li, W.X., 2006a. Geochemistry of the 755Ma Mundine Well dyke swarm, northwestern Australia: part of a Neoproterozoic mantle superplume beneath Rodinia? *Precambrian Research* 146, 1-15.
- Li, X.H., Li, Z.X., Sinclair, J.A., Li, W.X., Carter, G., 2006b. Revisiting the “Yanbian Terrane”: implications for Neoproterozoic tectonic evolution of the western Yangtze Block, South China. *Precambrian Research* 151, 14-30.
- Li, X.H., Liu, Y., Li, Q.L., Guo, C.H., Chamberlain, K.R., 2009. Precise determination of Phanerozoic zircon Pb/Pb age by multicollector SIMS without external standardization. *Geochemistry, Geophysics, Geosystems* 10(4), 1-21.
- Li, Y.H., Zhang, Z.J., Hou, K.J., Duan, C., Wan, D.F., Hu, G.Y., 2014. The genesis of Gongchanglin High-Grade-Iron Ores, Anshan-Benxi Area, Liaoning Province, NE China: Evidence from Fe-Si-O-S Isotopes. *Acta Geologica Sinica* 88(12), 2351-2372 (in Chinese with English abstract).
- Li, Z.H., Zhu, X.K., Tang, S.H., 2008. Characters of Fe isotopes and rare earth elements of banded iron formations from Anshan-Benxi area: implications for Fe source. *ACTA PETROLOGICA ET MINERALOGICA* 27(4), 285-290 (in Chinese with English abstract).
- Li, Z.H., Zhu, X.K., Tang, S.H., 2012. Fe isotope compositions of banded iron formation from Anshan-Benxi area: Constraints on the formation mechanism and Archean ocean environment. *Acta Petrologica Sinica* 28(11), 3545-3558 (in Chinese with English abstract).
- Li, Z.H., Zhu, X.K., Sun, J., 2014. Geochemical characters of Banded Iron Formation from Xinyu and North China. *Acta Petrologica Sinica* 30(5), 1279-1291 (in Chinese with English abstract).
- Li, Z.X., Bogdanova, S.V., Collins, A.S., Davidson, A., Waele, B.De, Ernst, R.E., Fitzsimons, I.C.W., Fuck, R.A., Gladkochub, D.P., Jacobs, J., Karlstrom, K.E., Lu, S., Natapov, L.M., Pease, V., Pisarevsky, S.A., Thrane, K., Vernikovsky, V., 2008. Assembly, configuration, and break-up history of Rodinia: A synthesis. *Precambrian Research* 160, 179-210.
- Lu, S.N., Li, H.K., Zhang, C.L., Niu, G.H., 2008. Geological and geochronological evidence for the Precambrian evolution of the Tarim Craton and surrounding continental fragments. *Precambrian Research* 160(1-2), 94-107.
- Ludwig, K.R., 2003. In: Ludwig, Kenneth R. (Ed.), *User's Manual for Isoplot 3.00: A Geochronological Toolkit for Microsoft Excel*.
- McLennan, S.M., 1989. Rare earth elements in sedimentary rocks: influence of provenance and sedimentary processes. *Rev. Mineral. Geochem.* 21, 169-200.
- Michard, A., Albarede, F., Michard, G., Minster, J.F., Charlou, J.L., 1983. Rare-earth elements and uranium in high-temperature solutions from East Pacific Rise hydrothermal vent field (13 N). *Nature* 303, 795-797.
- Michard, A., Michard, G., Stuben, D., Stoffers, P., Cheminée, J.L., Binard, N., 1993. Submarine thermal springs associated with young volcanoes: the Teahitia vents, Society islands Pacific Ocean. *Geochim. Cosmochim. Acta* 57, 4977-4986.
- Och, L.M., Shields-Zhou, G.A., 2012. The Neoproterozoic oxygenation event: Environmental perturbations and biogeochemical cycling. *Earth-Science Reviews* 110, 26-57.
- Planavsky, N., Bekker, A., Rouxel, O.J., Kamber, B., Hofmann, A., Knudsen, A., Lyons, T.W., 2010. Rare earth element and yttrium compositions of Archean and Paleoproterozoic Fe formations revisited: new perspectives on the significance and mechanisms of deposition. *Geochim. Cosmochim. Acta* 74, 6387-6405.
- Qi, L., D. Conrad Grégoire., 2000. Determination of Trace Elements in Twenty Six Chinese Geochemistry Reference Materials by Inductively Coupled Plasma-Mass Spectrometry. *The Journal of Geostandards and Geoanalysis* 24(1), 51-63.
- Rajabi, A., Canet, C., Rastad, E., Alfonso, P., 2015. Basin evolution and stratigraphic correlation of sedimentary-exhalative Zn-Pb deposits of the Early Cambrian Zarigan-Chahmir Basin, Central Iran. *Ore Geology Reviews* 64, 328-353.
- Rouxel, O., Bekker, A., Edwards, K., 2005. Iron isotope constraints on the Archean and Paleoproterozoic ocean redox state. *Science* 307, 1088-1091.
- Rouxel, O., Toner, B.M., Manganini, S.J., German, C.R., 2016. Geochemistry and iron isotope systematics of hydrothermal plume fall-out at East Pacific Rise 9°50'N. *Chemical Geology* 441, 212-234.
- Schröder, S., Grotzinger, J.P., 2007. Evidence for anoxia at the Ediacaran-Cambrian boundary: the record of redox-sensitive trace elements and rare earth elements in Oman. *Journal of the Geological Society, London* 164, 175-187.
- Shen, M.T., Xu, Z., Xu, M., 2018. Geological Geochemical and Fe Isotopic Compositions Characteristics and Its Implications on Xinyu Iron Deposit in Jiangxi Province. *Advances in Geosciences* 8(1), 158-172 (in Chinese with English abstract).
- Sial, A.N., Campos, M.S., Gaucher, C., Frei, R., Ferreira, V.P., Rielva, C., Nascimento, R.C., Pimentel, M.M., Pereira, N.S., Rodler, A., 2015. Algoma-type Neoproterozoic BIFs and related marbles in the Serido Belt (NE Brazil): REE, C, O, Cr and Sr isotope evidence. *J. S. Am. Earth Sci.* 61, 33-52.
- Sperling, E.A., Wolock, C.J., Morgan, A.S., Gill, B.C., Kunzmann, M., Halverson, G.P., Macdonald, F.A., Knoll, A.H., Johnston, D.T., 2015. Statistical analysis of iron geochemical data suggests limited late Proterozoic oxygenation. *Nature* 523, 451-454.
- Stern, R.J., Mukherjee, S.K., Nathan, R.M., Kamal, A., Peter, R.J., 2013. 750 Ma banded iron formation from the Arabian-Nubian Shield - Implications for understanding Neoproterozoic tectonics, volcanism, and climate

- change. *Precambr. Res.* 239, 79-94.
- Sun, S.S., McDonough, W.F., 1989. Chemical and isotopic systematics of oceanic basalts: Implications for mantle composition and processes. *Geological Society London Special Publications* 42(1), 313-345.
- Tang, S.H., Zhu, X.K., Li, J., Yan, B., Li, S.Z., Li, Z.H., Wang, Y., Sun, J., 2016. New Standard Solutions for Measurement of Iron, Copper and Zinc Isotope Compositions by Multi-collector Inductively Coupled Plasma-Mass Spectrometry. *ROCK AND MINERAL ANALYSIS* 35(2), 127-133 (in Chinese with English abstract).
- Wu, Y.B., Zheng, Y.F., 2004. A study on the genetic mineralogy of zircon and its restriction on the interpretation of U-Pb age. *CHINESE SCIENCE BULLETIN* 49(16), 1589 (in Chinese).
- Xinjiang Bureau of Geology and Mineral Resources (BGMR), 1993. *Regional Geology of Xinjiang Uygur Autonomous Region*. Geological Publishing House, Beijing, 8-33 (in Chinese).
- Xu, B., Xiao, S.H., Zou, H.B., Chen, Y., Li, Z.X., Song, B., Liu, D.Y., Zhou, C.M., Yuan, X.L., 2009. SHRIMP zircon U-Pb age constraints on Neoproterozoic Quruqtagh diamictites in NW China. *Precambrian Research* 168, 247-258.
- Xu, B., Zou, H.B., Chen, Y., He, J.Y., Wang, Y., 2013. The Sugetbrak basalts from northwestern Tarim Block of northwest China: Geochronology, geochemistry and implications for Rodinia breakup and ice age in the Late Neoproterozoic. *Precambrian Research* 236, 214-226.
- Xu, D.R., Wang, Z.L., Chen, H.Y., Hollings, Pete, Jansen, Nicholas, H., Zhang, Z.C., Wu, C.J., 2014. Petrography and geochemistry of the Shilu Fe-Co-Cu ore district, South China: Implications for the origin of a Neoproterozoic BIF system. *Ore Geology Reviews* 57, 322-350.
- Xu, Z.Q., He, B.Z., Zhang, C.L., Zhang, J.X., 2013. Tectonic framework and crustal evolution of the Precambrian basement of the Tarim Block in NW China: New geochronological evidence from deep drilling samples. *Precambrian Research* 235, 150-162.
- Yan, B., Zhu, X.K., Tang, S.H., Zhu, M.Y., 2010. Fe isotopic characteristics of the Neoproterozoic BIF in Guangxi Province and its Implications. *ACTA GEOLOGICA SINICA* 84(7), 1080-1086 (in Chinese with English abstract).
- Zhang, C.L., Ye, X.T., Zou, H.B., Chen, X.Y., 2016. Neoproterozoic sedimentary basin evolution in southwestern Tarim, NM China: New evidence from field observations, detrital zircon U-Pb ages and Hf isotope compositions. *Precambrian Research* 280, 31-45.
- Zhang, C.L., Zou, H.B., Li, H.K., Wang, H.Y., 2013. Tectonic framework and evolution of the Tarim Block in NW China. *Gondwana Research* 23, 1306-1315.
- Zhang, C.L., Zou, H.B., Santosh, M., Ye, X.T., Li, H.K., 2014. Is the precambrian basement of the Tarim Craton in NW china composed of discrete terranes?. *Precambrian Research* 254, 226-244.
- Zhang, J., Nozaki, Y., 1996. Rare earth elements and yttrium in seawater: ICP-MS determination in the East Caroline, Coral Sea, and South Fiji basins of the western South Pacific Ocean. *Geochim. Cosmochim. Acta* 60(23), 4631-4644.
- Zhou, X.Q., Chen, D.Z., Qing, H.R., Qian, Y.X., Wang, D., 2014. Submarine silica-rich hydrothermal activity during the earliest Cambrian in the Tarim Basin, Northwest China. *International Geology Review* 56(15), 1906-1918.
- Zhu, X.K., Guo, Y., Williams, R.J.P., O'Nions, R.K., Mattews, A., Belshaw, N.S., Canters, G.W., de Waal, E.C., Weser, U., Burgess, B.K., Salvato, B., 2002. Mass fractionation processes of transition metal isotopes, *Earth Planet Sci Lett* 200, 47-62.
- Zhu, X.K., Sun, J., Li, Z.H., 2019. Iron isotopic variations of the Cryogenian banded iron formations: A new model. *Precambrian Research* 331, 105359.

**Figure captions**




Fig. 1 (a) Sketch Precambrian geological map of Tarim Craton, showing the location of the Baijianshan BIF (modified after Lu et al., 2008). (b) Geological map of the Baijianshan Banded Iron Formation (Baijianshan BIF).




Fig. 2 Stratigraphical column of the Baijianshan BIF.




Fig. 3 Field photographs, hand specimen and mineral features of the Baijianshan BIF. (a) The tuff layers in the Baijianshan BIF. (b) Laminated Fe orebodies with local bend. (c) hand specimen. (d) Scanning electron microscopy (SEM) images. Detailed petrographic examination by SEM and transmitted light showing mineral assemblage: (e) Siderite. (f) Hematite. (g) Chlorite. (h) Sericite. Cht = Chert, Mt = Magnetite, Sd = Siderite, Q = Quartz, Hem = Hematite, Chl = Chlorite, Ser = Sericite.




Fig. 4 Concordia diagram of the zircon U-Pb ages of the tuff at the Baijianshan BIF.




Fig. 5 (a) Chondrite-normalized REE pattern of the basalts. Chondrite-normalized values after Sun and McDonough (1989). (b) PAAS-normalized REE+Y diagrams of BIFs from the Baijianshan BIF. The grayish zone represents the REY data from the Xinyu BIF (Li et al., 2014). PAAS-normalized values after McLennan (1989).




Fig. 6 (a) Bi-variation between Rb and Th of Baijianshan BIF samples; (b-i) Bi-variation between  $Al_2O_3$  and  $TiO_2$  (b),  $K_2O$  (c), Rb (d), V (e), Sc (f), Cs (g), Cr (h) and REE (i) of the Baijianshan BIF. Data for bulk-rock BIF samples are listed in supplementary Table 2.




Fig. 7  $\epsilon Nd(t)$  vs.  $(^{87}Sr/^{86}Sr)_i$  diagram of BIF and basalt samples in Baijianshan.




Fig. 8 (a)  $\epsilon_{\text{Nd}}(t = 0.737\text{Ga})$  vs.  $\text{Al}_2\text{O}_3$  diagram of the Baijianshan BIF; (b)  $\text{Ce}/\text{Ce}^*_{\text{PAAS}}$  vs.  $\text{Al}_2\text{O}_3$  diagram of the Baijianshan BIF.




Fig. 9 Sm/Yb vs. Eu/Sm ratios illustrating two component mixing model of the hydrothermal fluid of the Baijianshan BIF (after Alexander et al., 2008). BJS-Baijianshan BIF; XY-Xinyu BIF (Li et al., 2014); DHLT-Dahongliutan BIF (Hu et al., 2017). Data sets of Archean-Paleoproterozoic BIFs are from Planavsky et al. (2010). Average compositions of high-T ( $> 300\text{ }^\circ\text{C}$ ) hydrothermal fluids, low-T ( $< 200\text{ }^\circ\text{C}$ ) hydrothermal solutions and Pacific seawaters were after Bau and Dulski (1999), Michard et al. (1983) and Alibo and Nozaki (1999), respectively.




Fig. 10  $Pr/Pr^*_{PAAS}$  vs.  $Ce/Ce^*_{PAAS}$  diagram for the BIF at Baijianshan (after Bau and Dulski, 1996). The light blue area represents the data of Archean-Paleoproterozoic BIFs. Data sets of other BIFs are same as those of Fig. 9.




Fig. 11 Temporal variation in  $\delta^{57}Fe_{IRMM-014}$  of various ages BIFs in the world (Data of SW Greenland BIFs are from Dauphas et al.(2004); Manjeri Formation, Belingwe Belt, Zimbabwe BIFs are from Rouxel et al. (2005); Anshan-Benxi BIFs are from Li et al. (2012); Transvaal BIFs are from Johnson et al. (2003); Yuanjiacun BIFs are from Hou et al. (2014); Gongchangling BIFs from Li et al. (2014); Sanjiang BIF are from Yan et al. (2010) and Xinyu BIF are from Shen et al. (2008)).






Fig. 12 Cartoon model of the formation of the Baijianshan BIF (see details in the text).

### Appendix Table Captions

Appendix-Table 1 LA-ICP-MS U-Pb data for zircon from tuff in the first section of the Lapeiquan Formation

Appendix-Table 2 Geochemical compositions of the Baijianshan BIF

Appendix-Table 3 Sr-Nd isotopic data of the Baijianshan BIF

Appendix-Table 4 Fe isotopic data of the Baijianshan BIF

### Supplementary Table 1 Zircon U-Pb age data of the tuff from the Baijianshan BIF

Spot	U ppm	Th ppm	Th/U	<sup>206</sup> Pb/ <sup>238</sup> U Age ±1σ	<sup>207</sup> Pb/ <sup>235</sup> U Age ±1σ	<sup>207</sup> Pb/ <sup>206</sup> Pb	±1σ	<sup>207</sup> Pb/ <sup>235</sup> U U	±1σ	<sup>206</sup> Pb/ <sup>238</sup> U	±1σ		
2071-1	3079	831	0.27	1335	14	1360	21	0.0887	0.0012	2.8156	0.0427	0.2301	0.0025
2071-2	805	427	0.53	898	11	1154	20	0.1027	0.0014	2.1169	0.0369	0.1495	0.0018
2071-3	529	381	0.72	776	8	1009	18	0.0965	0.0016	1.7019	0.0307	0.1279	0.0014
2071-4	155	81	0.52	835	9	1350	22	0.1458	0.0022	2.778	0.0462	0.1382	0.0015
2071-5	184	101	0.55	751	8	1009	18	0.0999	0.0016	1.7022	0.0304	0.1236	0.0013
2071-6	297	252	0.85	1273	14	1318	20	0.0885	0.0012	2.6628	0.0414	0.2183	0.0023
2071-7	61	35	0.57	738	8	782	22	0.0693	0.0019	1.1591	0.0321	0.1212	0.0013
2071-8	3079	801	0.26	1365	15	1372	21	0.0881	0.0012	2.864	0.0432	0.2358	0.0026
2071-9	134	114	0.85	734	7	747	15	0.0653	0.0012	1.0863	0.0211	0.1206	0.0012
2071-10	40	25	0.64	740	8	676	30	0.0564	0.0025	0.9465	0.0419	0.1217	0.0013
2071-11	308	123	0.4	1122	13	1244	22	0.0917	0.0014	2.4035	0.0430	0.1902	0.0021
2071-12	239	187	0.78	738	8	744	12	0.0646	0.0009	1.0799	0.0170	0.1212	0.0012
2071-13	4616	2308	0.5	1283	14	1326	20	0.0887	0.0012	2.6928	0.0410	0.2202	0.0024
2071-14	104	151	0.69	735	9	778	24	0.0692	0.0021	1.1514	0.0362	0.1207	0.0015
2071-15	57	81	0.71	740	9	807	34	0.0724	0.0032	1.2148	0.0505	0.1217	0.0015
2071-16	239	131	0.55	792	8	1143	19	0.1155	0.0018	2.0821	0.0347	0.1307	0.0014
2071-17	356	185	0.52	776	8	976	18	0.0915	0.0016	1.615	0.0303	0.128	0.0014

2071-18	419	222	0.53	753	8	895	15	0.0828	0.0013	1.4142	0.0233	0.1238	0.0013
2071-19	149	69	0.46	741	8	817	26	0.0735	0.0022	1.2354	0.0387	0.1218	0.0013
2071-20	134	72	0.38	734	7	747	15	0.0653	0.0012	1.0863	0.0211	0.1206	0.0012
2071-21	525	242	0.46	741	8	777	13	0.0685	0.0010	1.1502	0.0191	0.1218	0.0013
2071-22	805	419	0.52	923	11	1167	20	0.1016	0.0014	2.1551	0.0374	0.1539	0.0018
2071-23	1066	661	0.62	748	8	795	12	0.0701	0.0010	1.1886	0.0184	0.123	0.0013
2071-24	435	318	0.73	1004	11	1194	22	0.0965	0.0015	2.2419	0.0408	0.1686	0.0018
2071-25	40	15	0.37	740	8	676	30	0.0564	0.0025	0.9465	0.0419	0.1217	0.0013
2071-26	200	96	0.48	787	8	1123	19	0.113	0.0017	2.0237	0.0339	0.1299	0.0014
2071-27	602	247	0.41	1081	12	1194	19	0.0891	0.0012	2.2408	0.0356	0.1825	0.0021
2071-28	88	165	0.49	733	9	748	27	0.0656	0.0023	1.0880	0.0392	0.1204	0.0015
2071-29	121	231	0.52	727	9	739	21	0.0650	0.0017	1.0699	0.0302	0.1194	0.0014
2071-30	3657	768	0.21	1188	13	1270	19	0.0893	0.0012	2.4939	0.0375	0.2025	0.0021
2071-31	185	67	0.36	737	7	740	13	0.0643	0.0010	1.0733	0.0182	0.121	0.0012
2071-32	61	22	0.54	738	8	782	22	0.0693	0.0019	1.1591	0.0321	0.1212	0.0013
2071-33	239	89	0.61	738	8	744	12	0.0646	0.0009	1.0799	0.0170	0.1212	0.0012

Supplementary Table 2 Geochemical compositions of Baijianshan BIF

Sample	BJS-1	BJS-2	BJS-3	BJS-4	BJS-5	BJS-6	BJS-7	BJS-8	BJS-9	BJS-10
Major elements (%)										
Fe <sub>2</sub> O <sub>3</sub>	53.60	41.52	56.64	52.26	30.01	36.85	41.01	45.22	39.78	48.55
SiO <sub>2</sub>	29.91	38.96	26.81	31.20	40.92	40.79	36.97	41.77	43.66	37.59
Al <sub>2</sub> O <sub>3</sub>	1.56	3.41	2.37	2.38	4.32	2.91	3.64	2.97	3.39	3.10
CaO	2.48	0.54	1.10	0.34	3.63	2.05	0.76	0.23	0.47	0.51
MgO	0.87	1.50	1.69	1.29	2.62	1.86	1.48	1.29	0.80	0.84
TiO <sub>2</sub>	0.23	0.41	0.24	0.25	0.44	0.27	0.37	0.40	0.44	0.32
MnO	1.10	0.04	0.40	0.09	0.19	0.09	0.04	0.00	0.01	0.00
P <sub>2</sub> O <sub>5</sub>	1.47	0.29	0.41	0.16	1.58	0.35	0.15	0.32	0.47	0.46
K <sub>2</sub> O	0.13	0.06	0.09	1.02	1.33	0.90	0.41	1.25	1.27	1.39
Na <sub>2</sub> O	0.16	0.06	0.10	0.13	1.42	0.54	0.42	0.87	1.12	1.15
LOI	9.21	3.69	3.75	1.93	4.42	4.11	3.04	1.55	1.37	1.05
FeO	<0.02	9.47	6.49	8.89	9.90	9.46	11.64	4.39	7.66	5.28
Total	100.74	99.95	100.09	99.94	100.78	100.18	99.93	100.26	100.44	100.24
Trace elements (ppm)										
Cr	32.5	63.2	50.6	42.5	83.5	47.7	45.6	80.6	68.8	15.6
Ni	10.3	11.2	9.67	8.83	14.0	10.0	11.2	14.0	19.3	10.6
Co	64.7	32.5	34.2	37.6	21.3	33.1	17.6	100	141	85.2
Sc	6.25	6.35	7.91	7.78	8.19	6.81	6.82	7.60	7.51	6.81
Ba	147	25.4	94.6	1505	527	647	232	129	277	196
Sr	243	43.9	88.9	32.0	170	81.0	20.3	78.8	73.7	37.1
Rb	3.71	2.84	7.75	47.8	85.2	41.4	17.2	79.3	17.5	34.4
Zr	42.1	29.4	64.1	68.0	67.2	61.9	77.8	60.8	68.4	65.7
Hf	0.98	0.69	1.14	1.27	2.00	1.45	1.78	1.49	1.71	1.42
Ta	0.27	0.42	0.31	0.33	0.56	0.35	0.32	0.57	0.70	0.48
Ga	2.88	5.59	4.15	3.90	5.97	4.61	6.15	5.42	5.80	4.25
Cu	11.4	15.0	13.5	8.87	18.3	11.3	9.06	8.03	9.09	5.66
Zn	34.0	38.7	25.0	30.0	33.6	30.2	46.1	30.6	28.9	17.6
Pb	6.84	5.86	3.09	15.80	7.79	5.22	4.17	5.60	6.15	5.77
U	7.33	0.57	1.25	0.44	0.63	0.34	0.66	0.39	0.31	0.26
Th	1.92	3.26	2.73	2.31	4.84	2.65	3.20	3.48	3.79	2.64
Nb	3.02	5.66	4.47	4.80	5.75	4.84	5.81	4.91	6.34	5.76
La	9.77	15.9	16.3	11.4	18.5	6.45	5.64	15.3	19.6	8.38
Ce	22.5	35.9	38.0	26.9	40.7	15.0	13.3	35.4	44.6	19.0
Pr	2.82	4.39	4.69	3.22	5.10	1.83	1.65	4.00	5.28	2.30
Nd	12.9	19.0	20.2	13.5	22.2	7.91	7.06	16.5	22.2	9.67
Sm	3.11	4.22	4.51	2.92	5.06	1.92	1.72	3.38	4.60	2.15
Eu	0.81	1.02	1.08	0.68	1.25	0.47	0.39	0.77	1.11	0.51
Gd	3.74	4.19	4.65	3.04	5.63	2.23	1.86	3.39	4.41	2.31
Tb	0.62	0.66	0.76	0.51	0.90	0.39	0.31	0.55	0.68	0.41
Dy	4.37	4.01	5.05	3.49	6.00	2.71	2.14	3.63	4.24	2.86
Y	32.7	24.3	31.7	21.4	38.5	16.3	11.4	21.7	24.0	17.3
Ho	1.03	0.87	1.14	0.81	1.34	0.63	0.48	0.83	0.92	0.65
Er	3.01	2.34	3.30	2.43	3.77	1.83	1.39	2.47	2.59	1.93
Tm	0.46	0.35	0.50	0.39	0.55	0.29	0.23	0.39	0.39	0.30
Yb	3.09	2.19	3.37	2.71	3.65	2.01	1.67	2.70	2.66	2.12
Lu	0.50	0.39	0.54	0.45	0.58	0.33	0.27	0.45	0.42	0.34
∑REE	68.73	95.43	104.09	72.44	115.23	43.99	38.11	89.76	113.7	52.94
Y/Ho	31.75	28.00	27.81	26.32	28.73	26.04	23.85	26.08	26.23	26.57
La/La*	1.30	1.12	1.06	1.00	1.14	1.09	1.02	1.04	1.06	1.04
Ce/Ce*	0.98	0.99	1.00	1.02	0.96	1.00	1.00	1.04	1.01	0.99
Y/Y*	1.23	1.04	1.06	1.02	1.09	1.00	0.90	1.00	0.98	1.02
Pr/Pr*	0.96	0.98	0.99	0.99	0.99	0.98	1.00	0.97	0.98	0.99
Eu/Eu*	1.10	1.14	1.10	1.06	1.09	1.06	1.01	1.06	1.16	1.08
La <sub>N</sub> /Yb <sub>N</sub>	0.23	0.54	0.36	0.31	0.37	0.24	0.25	0.42	0.54	0.29

Sample	BJS-12	BJS-13	BJS-14	BJS-15	BJS-16	BJS-17	BJS-18	2702H1	2702H2	2702H3
--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------

Major elements (%)

TFe <sub>2</sub> O <sub>3</sub>	21.79	28.86	44.82	42.63	26.98	13.11	40.45	11.61	11.70	11.98
SiO <sub>2</sub>	55.06	49.87	41.24	43.65	53.64	59.90	42.80	39.94	40.91	44.55
Al <sub>2</sub> O <sub>3</sub>	5.24	4.93	2.81	2.86	4.83	7.58	3.84	11.36	11.45	10.76
CaO	0.81	1.02	0.75	0.75	0.76	0.84	0.65	9.77	9.31	7.45
MgO	1.57	1.51	0.74	0.86	1.36	2.32	1.31	8.40	8.54	10.38
TiO <sub>2</sub>	0.60	0.53	0.35	0.25	0.59	0.86	0.41	3.66	3.80	3.60
MnO	0.03	0.04	0.05	0.03	0.03	0.02	0.03	0.24	0.25	0.23
P <sub>2</sub> O <sub>5</sub>	0.20	0.35	0.49	0.40	0.23	0.19	0.54	0.46	0.43	0.38
K <sub>2</sub> O	3.11	2.32	0.26	0.17	1.86	3.78	1.98	1.35	1.27	0.36
Na <sub>2</sub> O	0.94	1.25	0.14	0.19	0.63	0.77	0.88	0.94	0.98	0.59
LOI	2.06	1.97	4.57	2.89	2.9	3.19	1.92	12.02	11.39	9.60
FeO	7.97	7.26	3.89	5.29	6.06	6.57	5.31	-	-	-
Total	99.38	99.91	100.11	99.97	99.87	99.13	100.12	99.75	100.03	99.88
Trace elements (ppm)										
Cr	118	81.0	59.7	10.5	86.2	112	50.9	214	214	306
Ni	15.6	14.5	12.1	11.0	15.6	12.3	12.6	162	166	217
Co	76.1	70.2	33.7	34.7	54.0	27.1	45.3	48.0	62.9	50.7
Sc	12.2	14.4	7.97	8.11	11.7	15.1	8.11	26.4	26.1	25.7
Ba	662	562	111	19.2	359	1219	385	877	864	296
Sr	45.0	53.3	45.8	26.1	32.6	37.0	48.9	205	209	150
Rb	111	95.6	7.36	6.13	113	184	85.6	34.4	32.2	12.3
Zr	101	93.4	60.0	87.9	79.9	116	79.7	259	251	228
Hf	2.63	2.37	1.23	1.50	2.11	2.99	1.92	5.35	5.74	6.00
Ta	0.88	0.75	0.37	0.36	0.64	0.84	0.53	2.24	2.33	2.14
Ga	7.95	7.63	4.39	4.63	7.12	10.3	5.97	18.3	17.2	18.8
Cu	27.6	21.6	8.12	3.56	21.4	36.0	12.4	-	-	-
Zn	62.1	40.0	26.0	23.8	35.3	37.0	29.6	-	-	-
Pb	7.80	8.24	3.54	5.32	7.64	15.4	5.75	-	-	-
U	0.47	0.69	0.39	0.40	0.72	0.82	0.37	1.45	1.09	0.92
Th	7.26	5.14	2.81	2.97	5.52	7.11	4.11	3.32	3.37	3.18
Nb	8.35	8.21	4.19	4.86	6.80	9.50	6.11	38.0	36.5	33.6
La	26.8	20.3	13.0	9.82	19.1	22.5	17.3	36.9	34.1	33.7
Ce	58.3	44.2	30.2	22.8	42.0	46.1	39.0	77.2	72.9	74.9
Pr	7.03	5.35	3.73	2.77	5.09	5.51	4.81	9.37	9.15	9.28
Nd	29.3	22.4	16.2	11.7	21.3	22.6	20.7	37.9	36.6	40.0
Sm	6.46	4.95	3.67	2.65	4.70	4.96	4.70	8.06	8.20	9.11
Eu	1.33	1.15	0.91	0.66	1.06	1.01	1.12	1.41	1.85	2.91
Gd	6.39	5.03	4.02	3.03	4.64	4.93	5.01	7.69	7.82	8.87
Tb	1.00	0.80	0.67	0.52	0.72	0.79	0.82	1.21	1.22	1.35
Dy	5.98	4.98	4.53	3.71	4.26	4.84	5.36	6.57	7.04	7.04
Y	28.5	24.9	27.9	22.0	20.1	26.1	31.9	34.8	34.0	32.1
Ho	1.20	1.04	1.02	0.85	0.86	1.01	1.17	1.25	1.29	1.31
Er	3.23	2.94	3.00	2.59	2.33	2.82	3.38	3.11	3.15	3.20
Tm	0.45	0.43	0.46	0.40	0.34	0.41	0.50	0.39	0.40	0.39
Yb	2.87	2.93	3.06	2.76	2.20	2.69	3.36	2.28	2.36	2.28
Lu	0.43	0.45	0.49	0.45	0.33	0.41	0.53	0.34	0.32	0.32
∑REE	150.77	116.95	84.95	64.71	108.93	120.58	107.75	193.68	186.4	194.67
Y/Ho	23.75	23.94	27.35	25.76	23.29	25.84	27.26	27.84	26.36	24.50
La/La*	1.06	1.07	1.09	1.02	1.06	1.09	1.10	1.14	1.06	1.14
Ce/Ce*	0.98	0.98	0.99	1.00	0.98	0.95	0.98	0.98	0.98	1.01
Y/Y*	0.85	0.88	1.04	0.99	0.84	0.95	1.02	0.89	0.83	0.78
Pr/Pr*	1.00	1.00	0.99	0.99	1.00	1.00	0.99	0.97	0.99	0.95
Eu/Eu*	0.97	1.08	1.10	1.08	1.07	0.96	1.08	0.54	0.69	0.97
La <sub>N</sub> /Yb <sub>N</sub>	0.69	0.51	0.31	0.26	0.64	0.62	0.38	10.99	9.82	10.04

Supplementary Table 3 Sr-Nd isotopic data of the Baijianshan BIF

Sample	<sup>87</sup> Rb/ <sup>86</sup> Sr	<sup>87</sup> Sr/ <sup>86</sup> Sr	2δ	( <sup>87</sup> Sr/ <sup>86</sup> Sr) <sub>i</sub>	<sup>147</sup> Sm/ <sup>144</sup> Nd	<sup>143</sup> Nd/ <sup>144</sup> Nd	2δ	T <sub>DM</sub> (Ga)	εNd(t)
BJS-1	0.0433	0.717927	0.000004	0.7175	0.1457	0.512425	0.000006	1.63	0.65
BJS-4	4.2608	0.727751	0.000009	0.6829	0.1308	0.512394	0.000003	1.39	1.46
BJS-5	1.4305	0.728893	0.000008	0.7138	0.1378	0.512363	0.000003	1.58	0.19
BJS-7	2.4192	0.729423	0.000009	0.7040	0.1473	0.512376	0.000002	1.78	-0.45
BJS-8	2.8555	0.718681	0.000003	0.6886	0.1238	0.512351	0.000004	1.36	1.27
BJS-10	2.6390	0.723954	0.000005	0.6962	0.1344	0.512364	0.000002	1.51	0.53
BJS-12	7.0891	0.740795	0.000007	0.6662	0.1333	0.512311	0.000003	1.59	-0.40
BJS-16	9.9936	0.746329	0.000008	0.6412	0.1334	0.512317	0.000003	1.58	-0.30
BJS-18	5.0054	0.731954	0.000008	0.6793	0.1373	0.51237	0.000004	1.56	0.38
2702H1	0.4737	0.711034	0.000009	0.7061	0.1286	0.512662	0.000003	0.88	6.90
2702H2	0.4355	0.711156	0.000009	0.7066	0.1354	0.512694	0.000006	0.89	6.88
2702H3	0.2322	0.710822	0.000012	0.7084	0.1377	0.512644	0.000005	1.02	5.69

Supplementary Table 4 Fe isotopic data of the Baijianshan BIF

Sample	δ <sup>57</sup> Fe <sub>IRMM-014</sub>	2σ	δ <sup>56</sup> Fe <sub>IRMM-014</sub>	2σ
--------	--	----	--	----

BJS-2	2.10	0.09	1.46	1.46
BJS-5	1.78	0.02	1.24	1.24
BJS-7	2.76	0.09	1.89	1.89
BJS-9	2.49	0.05	1.69	1.69
BJS-12	3.05	0.04	2.17	2.17
BJS-17	2.80	0.01	1.91	1.91